

Prevalence of Generalized Anxiety Disorder and Associated Risk Factors Among Medical Students in Sudan: A Cross-Sectional Study at Omdurman Islamic University

Khalid Osman Mohamed,¹ Ahmed A Lemam Ahmed,² ElShimaa Ammar Zaki,³ Sozan Mudather Soumit,⁴ Wamda Ahmed Ali,⁵ Asmaa Mohamed Abbas.⁶

Abstract

Background: Generalized Anxiety Disorder (GAD) is a mental illness that significantly affects various domains of daily functioning. Limited research has been conducted on GAD among medical students in Sudan, particularly during the socio-political and economic crises. This study aimed to assess the prevalence of GAD, identify risk factors, and evaluate its impact on academic performance and daily activities among Sudanese medical students. **Methods:** A cross-sectional study was conducted among undergraduate medical students at Omdurman Islamic University. Data were collected using a self-administered online questionnaire via Google Forms, consisting of two parts: socio-demographic information and the Generalized Anxiety Disorder Questionnaire (GAD-7), a validated tool for screening and measuring the severity of GAD. **Results:** A total of 374 medical students participated, with 64.7% being female. The GAD-7 scores were high (above 9), suggesting GAD among 33.7% of participants, with severity levels of 41.2% for mild anxiety, 21.4% for moderate anxiety, and 12.3% for severe anxiety. Comparison of means showed significant associations between GAD and female students ($p < 0.001$) and students with chronic diseases ($p = 0.034$). GAD significantly impacted daily activities ($p < 0.001$). Multiple logistic regression analysis found that students in the final year had significantly higher GAD-7 scores (Adjusted Odds Ratio=4.25, 95% Confidence Interval=1.27-14.22). **Conclusions:** The higher scores on the GAD-7 measure among Sudanese medical students are concerning. This emphasizes the urgent need to raise awareness, normalize mental health discussions, and provide accessible counseling services tailored to the students' needs.

Introduction

Mental disorders are the leading cause of disability worldwide. Individuals with significant mental health issues tend to have a lifespan that is 10 to 20 years shorter than that of the average person.¹

Generalized Anxiety Disorder (GAD) is a mental condition characterized by persistent and excessive anxiety and worry about various events or activities, such as school or work performance. These symptoms occur on most days for at least six months and can hinder functioning in social, occupational, or other domains.²

The disease is highly prevalent; in the United States, it is estimated that 6.8 million adults have GAD, with only 43.2% receiving treatment.³ It is also estimated that 5.7% of U.S. adults experience generalized anxiety disorder at some point in their lives.⁴

Studies have consistently shown that females are more likely than males to develop GAD,^{5,6} with the prevalence being twice as high for them.⁷ There are other risk factors associated with GAD, such as genetic factors,⁸ and chronic diseases like diabetes mellitus,⁹ asthma,¹⁰ and systemic lupus erythematosus.¹¹

The demanding nature of medical school with challenging training programs, both academically and emotionally across all professions, places medical students at a higher risk for GAD compared to the general population. Studies have revealed that 29% to 38% of medical students experience GAD, in contrast to the 3% to 25% prevalence observed in the general population.¹² The academic years in medical school are filled with numerous challenges, including demanding coursework, difficult exams, and extensive study hours.

Studies have shown that the prevalence of GAD among medical students varies between countries. A study conducted in the USA reported that 65.9% of medical students exhibited symptoms of anxiety.¹³ Meanwhile, in Saudi Arabia, 69% of medical students were found to have varying degrees of GAD,¹⁴ and in Egypt, the rate was a bit higher at 77.1%.¹⁵

The effects of GAD on medical students are profound. A study conducted in Mexico aimed to assess the impact of GAD on university students during the COVID-19 pandemic on academic performance. The findings revealed a significant negative effect of anxiety on students' academic performance.¹⁶ Similarly, a study

¹ MBBS. University of Bahri, Khartoum, Sudan.

² MBBS. Gadarif Teaching Hospital, Gadarif, Sudan.

³ Fifth-year Medical Student. Omdurman Islamic University, Omdurman, Sudan.

⁴ MBBS. Police Teaching Hospital, Khartoum, Sudan.

⁵ Third-year Pharmacy Student. Omdurman Islamic University, Khartoum, Sudan.

⁶ MBBS. University of Al Fashir, Al-Fashir, Sudan.

About the Author: ElShimaa Ammar Zaki is a fifth-year medical student currently studying at Omdurman Islamic University in Omdurman, Sudan.

Correspondence:

Khalid Osman Mohamed.

Address: QH3W+24W، الخرطوم بحري-الكلدو جنوب، السودان، Alkadroo, Sudan

Email: Khaledothman94@gmail.com

Editor: Francisco J. Bonilla-Escobar

Student Editors: Marcel Chee &

Diego Carrion Alvarez

Proofreader: Amy Phelan

Submission: May 24, 2023

Revisions: Aug 7, 2023; Nov 17, 2023

Responses: Aug 15, 2023; Feb 5, 2024

Acceptance: Mar 22, 2024

Publication: Mar 25, 2024

Process: Peer-reviewed

among medical students in Syria demonstrated a negative association between anxiety and academic performance.¹⁷

Anxiety not only affects academic performance but also impairs the ability to achieve work goals, manage household tasks, and interact with others. Several studies have reported that GAD is associated with a poor quality of life.^{18,19}

While the global prevalence rates are alarming, the situation in Sudan presents unique challenges; in 2020, a study among medical students revealed prevalence rates of depression at 75%, anxiety at 55.3%, and stress at 51.8%.²⁰ In recent years, Sudan has faced political and environmental instability, significantly impacting the mental health of its population. Medical students in Sudan are particularly struggling to manage the requirements of their academic programs with the unstable socio-political environment, potentially exacerbating their mental health issues.²¹

Given the significant global and local implications of GAD among medical students, particularly amidst the critical socio-political challenges in Sudan, there is a pressing need for more comprehensive studies within the country. This study aimed to determine the prevalence of GAD, explore associated risk factors, and evaluate its impact on academic performance and daily activities among medical students at Omdurman Islamic University to contribute valuable insights that can guide more effective mental health support strategies for medical students in Sudan.

Methods

Ethical Considerations

This study received ethical approval on July 21, 2022, from the Ethics Committee of the Faculty of Medicine at Omdurman Islamic University (Ethical Approval No. 3/2022). All participants provided informed consent online.

Study Setting and Design

This descriptive cross-sectional institution-based study was conducted among medical students at Omdurman Islamic University from October to December 2022. The university, located in Omdurman City, Khartoum State, Sudan, had approximately 26,000 students in 2022.

Sampling Strategy

The sample size was calculated to be 379 using the following formula: $n = Z^2 p[1-p]/d^2$, where n = calculated sample size, p is the prevalence of anxiety based on a previous study = 0.553, Z is the confidence level used = 1.96, and d = the level of precision (0.05).²⁰

Participants for the study were selected using a systematic random sampling technique. The student list was obtained from the faculty administration. To determine the sampling interval, we divided the total number of medical students by the calculated sample size, resulting in an interval of five. A random number

generator was used to select the first study participant from the first five students on the list. Subsequently, every fifth student from this initial point was selected to participate in the study. The inclusion criteria included medical students actively enrolled at the university. There were no exclusion criteria. At the start of the questionnaire, participants were informed about the purpose of the study, and assured that their involvement was entirely voluntary and that their anonymity would be preserved. Only those who gave their consent were able to fill out the questionnaire.

Data Collection Tools

Data were collected using a self-administered online questionnaire via Google Forms. The selected participants received the questionnaire link through WhatsApp. The questionnaire consisted of two sections: The first part was socio-demographic information, specific risk factors, and academic performance data: (gender, age, marital status, academic year, residence, living situation, grade point average (GPA), and medical history). The second part assessed GAD severity using the Generalized Anxiety Disorder Questionnaire (GAD-7).

Academic Performance

The academic performance of students was evaluated using their GPA, as per the system of the faculty of Medicine at Omdurman Islamic University administration. GPAs were categorized on a scale from 0 to 4, with the following classifications: Distinction (3.5-4), Very Good (3-3.49), Good (2.50-2.99), Pass (2-2.49), and Fail (<2).

Generalized Anxiety Disorder (GAD-7)

The GAD-7 is a valid and efficient tool for screening GAD and its severities. Scores are interpreted as follows: <5 (normal), 5-9 (mild anxiety), 10-14 (moderate anxiety), and 15-21 (severe anxiety). A cut-off score of 10 identifies cases of GAD, with a sensitivity of 89% and a specificity of 82%.^{22,23}

Statistical Analysis

Data were analyzed using SPSS version 28. Descriptive statistics, such as frequencies, means, standard deviations, and percentages, were employed to describe the dataset. A GAD-7 score of 10 or more was considered indicative of high anxiety levels. The relationship between risk factors and GAD was analyzed using t-tests and one-way ANOVA. Statistical significance was set at $P = 0.05$ or less. To measure the effect size, Cohen's d was used for t-tests, and eta-squared was used for one-way ANOVA tests. Simple and multiple logistic regression were performed to predict Generalized Anxiety Disorder from various independent variables.

Results

Sociodemographic and Risk Factor Profiles

A total of 374 medical students participated in this study. Most of them were female (64.7%). Their mean age was 21.07 ± 2.18 , ranging from 16 to 29 years. Socio-demographic data are presented in [Table 1](#).

Table 1. Sociodemographic and Risk Factor of Medical Students at Omdurman Islamic University, n=374.

| Variable | n | % |
|------------------------------------|-----|------|
| Gender | | |
| Male | 132 | 35.3 |
| Female | 242 | 64.7 |
| Academic year | | |
| 1st year | 80 | 21.4 |
| 2nd year | 85 | 22.7 |
| 3rd year | 69 | 18.8 |
| 4th year | 79 | 21.1 |
| 5th year | 61 | 16.3 |
| Residency city | | |
| Khartoum | 82 | 21.9 |
| Bahri | 42 | 11.2 |
| Omdurman | 247 | 66.0 |
| Madani | 3 | 0.8 |
| Marital status | | |
| Single | 366 | 97.9 |
| Married | 8 | 2.1 |
| Residency situation | | |
| First degree family | 219 | 58.6 |
| 2nd degree family | 27 | 7.2 |
| Dormitory | 108 | 28.8 |
| In apartments with other students | 17 | 4.5 |
| With husband | 1 | 0.3 |
| Alone | 2 | 0.6 |
| Chronic Disease Presence | | |
| yes | 37 | 9.9 |
| No | 337 | 90.1 |
| Exercise Frequency Per Week | | |
| No | 299 | 79.9 |
| 1-2 times | 21 | 5.6 |
| 3-4 times | 29 | 7.8 |
| 5 times | 17 | 4.5 |
| More than 5 times | 8 | 2.1 |
| Smoking status | | |
| Ex-smoker | 7 | 1.9 |
| Smoker | 8 | 2.1 |
| Not smoker | 359 | 96.0 |

Out of the 374 medical students who participated in the study, 101 of them reported engaging in regular exercise. It was found that nearly 50% of these students reported engaging in regular cardio exercises, while approximately 40% stated that they regularly engage in walking exercises, as depicted in [Figure 1](#).

Prevalence and Severity of Generalized Anxiety Disorder among Medical Students at Omdurman Islamic University

According to the GAD-7 scale, 33.7% of the medical students exhibited high GAD scores, indicating scores of 10 or more. The severity of these anxiety levels is detailed in [Figure 2](#).

Figure 1. Types of Exercises Performed by Participants, Sudan, 2022, n=101.





Figure 2. Prevalence and Severity of Generalized Anxiety Disorder (GAD) among Participants at Omdurman Islamic University, n=374.



Assessment of the Risk Factors for GAD among Medical Students at Omdurman Islamic University

A comparison of means was conducted to examine variable differences in GAD-7 scores. The results indicated that Female medical students (M = 7.80) had significantly higher GAD-7 scores than male medical students (M = 6.06) (p < 0.001). Additionally, no significant differences were found between age, academic year, residence city, marital status, residence situation, smoking status, and GAD-7 scores (all p-values > 0.05). However, medical students with chronic diseases had significantly higher GAD-7 scores (M = 8.64) than those without (M = 7.03) (p = 0.034). Furthermore, there was a significant difference in the GAD-7 scores between medical students who reported performing regular exercise (M = 7.41) and those who did not (M = 6.29, p = 0.05. [Table 2](#)).

Table 2. Mean Scores, Standard Deviations, and Effect Sizes of GAD-7 Scores Across Socio-Demographic and Risk Factors.

| Variable | Mean | Standard deviation | p value | Effect size |
|-----------------------------------|-------|--------------------|---------|-------------|
| Gender | | | | |
| Male | 6.07 | 4.40 | <0.001 | -0.40 |
| Female | 7.81 | 4.30 | | |
| Age | | | | |
| Less than 18 years | 7.14 | 5.14 | 0.36 | 0.009 |
| 18 to 20 years | 6.96 | 4.47 | | |
| 21 to 25 years | 7.47 | 4.36 | | |
| More than 25 years | 5.00 | 3.74 | | |
| Academic year | | | | |
| 1st year | 6.86 | 4.63 | 0.35 | 0.01 |
| 2nd year | 6.62 | 4.50 | | |
| 3rd year | 7.19 | 3.81 | | |
| 4th year | 7.52 | 4.70 | | |
| 5th year | 8.02 | 4.21 | | |
| Residency city | | | | |
| Khartoum | 7.40 | 4.54 | 0.85 | 0.002 |
| Bahri | 7.40 | 3.62 | | |
| Omdurman | 7.07 | 4.51 | | |
| Madani | 8.66 | 2.31 | | |
| Marital status | | | | |
| Single | 7.14 | 4.36 | 0.11 | -0.56 |
| Married | 9.62 | 6.19 | | |
| Residency situation | | | | |
| First degree family | 7.18 | 4.49 | 0.05 | 0.02 |
| 2nd degree family | 5.30 | 4.40 | | |
| Dormitory | 7.49 | 4.15 | | |
| In apartments with other students | 8.00 | 4.32 | | |
| With husband | 15.00 | N/A | | |
| Chronic Disease | | | | |
| Presence | | | | |
| yes | 8.65 | 5.57 | 0.03 | 0.37 |
| No | 7.04 | 4.24 | | |
| Exercise | | | | |
| Yes | 7.42 | 4.47 | 0.05 | 0.25 |
| No | 6.30 | 4.07 | | |
| Smoking status | | | | |
| Ex-smoker | 9.14 | 4.67 | 0.45 | 0.004 |
| Smoker | 7.87 | 6.08 | | |
| Not smoker | 7.14 | 4.37 | | |

A multiple logistic regression analysis was conducted to investigate the relationship between several predictor variables and the presence of Generalized anxiety disorder. The results show that being female ($p < 0.001$, AOR = 2.399) and being in the 5th academic year ($p = 0.019$, AOR = 4.246) were associated with higher odds of anxiety ([Table 3](#)).

Impact of GAD on academic performance among medical students at Omdurman Islamic University

The mean GPA of the medical students is 3.25 ± 0.525 SD. To evaluate the impact of Generalized Anxiety Disorder on students' academic performance, we compared the means of the GPA across various GAD severity levels using an ANOVA test. The findings, detailed in [Table 4](#), show no significant differences in the GPAs across the different levels of anxiety among the students.

Table 3. Multiple Logistic Regression of Predictors of Generalized Anxiety Disorder (GAD).

| Predictor Variable | Adjusted Odds Ratio (AOR) | 95% CI | p-value |
|--------------------|---------------------------|-------------|---------|
| Age | 0.90 | 0.76, 1.06 | 0.21 |
| Gender | | | |
| Male | (Reference) | - | - |
| Female | 2.40 | 1.46, 3.95 | <0.001 |
| Academic year | | | |
| 1st year | (Reference) | - | - |
| 2nd year | 1.30 | 0.65, 2.60 | 0.46 |
| 3rd year | 2.03 | 0.89, 4.63 | 0.09 |
| 4th year | 2.34 | 0.91, 6.01 | 0.08 |
| 5th year | 4.25 | 1.27, 14.22 | 0.02 |
| Marital status | | | |
| Not married | (Reference) | - | - |
| Married | 0.66 | 0.12, 3.52 | 0.62 |
| Chronic disease | | | |
| No | (Reference) | - | - |
| Yes | 0.78 | 0.36, 1.67 | 0.52 |
| Exercise | | | |
| No | (Reference) | - | - |
| Yes | 1.20 | 0.68, 2.14 | 0.53 |

Table 4. Mean Scores and Standard Deviations of grade point average (GPA) Across Generalized Anxiety Disorder (GAD) Severities.

| GAD severities | GPA Mean | Standard deviation | p-value |
|------------------|----------|--------------------|---------|
| No Anxiety | 3.23 | 0.623 | 0.160 |
| Mild Anxiety | 3.32 | 0.469 | |
| Moderate Anxiety | 3.16 | 0.512 | |
| Severe Anxiety | 3.26 | 0.491 | |

Table 5. The Impact of Generalized Anxiety Disorder (GAD) Severity on Daily Life Activities.

| If you checked any problems, how difficult have they made it for you to do your work, take care of things at home, or get along with other people? | Mean | Standard Deviation | p-value | Effect Size |
|--|----------------------|--------------------|---------|-------------|
| | Not difficult at all | 4.11 | 3.67 | <0.001 |
| Somewhat difficult | 6.86 | 3.25 | | |
| Very difficult | 11.95 | 3.95 | | |
| Extremely difficult | 11.81 | 3.60 | | |

Impact of GAD on daily life activities among medical students at Omdurman Islamic University

A comparison of the means was conducted to study the impact of GAD on daily activities among students. Significantly higher mean GAD-7 scores were found among students who reported extremely difficult (11.80) and very difficult (11.95) daily activities, and lower mean GAD-7 scores were found among students who reported not difficult (4.10) and somewhat difficult (6.86) daily activities ($p < 0.001$, [Table 5](#)).

Discussion

Our study found that 33.7% of medical students had significantly high GAD-7 scores (scores > 9), suggesting the possibility of GAD. This rate is slightly higher than the 29% recently reported among medical students at the University of Khartoum, Sudan.²⁴ Another study, conducted during the COVID-19 lockdown in Sudan reported a prevalence rate of 55.3%; however, this study employed the DASS-21 scale, a different assessment tool, potentially explaining the variance in rates.²⁰

Participants in our study were exposed to a complex social and political environment characterized by political unrest and economic hardship. Even following the COVID-19 pandemic, colleges often faced closures due to the unstable political situation. These factors could heighten anxiety and uncertainty about future careers among students, thereby impacting their mental well-being.

Comparatively, studies in Saudi Arabia and the USA reported prevalence rates of 31.2% and 30.6%, respectively.^{13,25} In contrast, studies in Egypt reported higher anxiety rates of 77.1% and 56%; however, these studies utilized the DASS-42 tool, which might account for the higher rates.^{26,27}

A high level of stress often accompanies medical school. A comparative study assessed stress levels among students from various disciplines. The findings revealed that 54.3% of medical students reported experiencing significant stress, compared to 36.6% of arts students, 32% of business students, and 15.3% of engineering students.²⁸

In our study, a significant association was observed between female medical students and GAD ($p < 0.001$). This finding is consistent with studies conducted among medical students in several countries, including Saudi Arabia,²⁹ Turkey,³⁰ Egypt,³¹ and the United States.¹³ Some studies suggest that specific biological factors, such as abnormalities in female hormones, might contribute to increased anxiety in women, potentially leading to a higher risk of GAD compared to men.^{32,33} Another study indicated that female students may be more susceptible to GAD due to their perception of patient contacts and autopsy-related duties as stressful, with exams also being a significant source of stress.³⁴ Moreover, multiple studies have reported that female medical students are more likely to experience of imposter syndrome, which is strongly associated with anxiety and other psychological conditions.^{35,36}

The study revealed that final-year medical students were more likely to experience generalized anxiety disorder (GAD) than students in earlier academic years (AOR= 4.25) ($p=0.019$). This can be explained by the increase in academic and practical workloads, as well as the stress associated with final exams, during the last year. This aligns with previous research indicating a progressive increase in stress throughout medical education.³⁷ Existing studies suggest that mental health often declines upon

entering medical school and remains challenging throughout the program.³⁸

Regarding academic performance, our study did not find a significant difference in GPA scores across different GAD severities. This finding contradicts the results of many studies, including a meta-analysis of 238 studies, which found a significant association between increased GAD severity and poorer academic performance.³⁹ However, our results align with other research that found no association between academic performance and GAD.⁴⁰ Some studies have even suggested that anxiety levels might increase with higher GPA scores, as students may become more concerned about maintaining their grades.²⁵ These variations could be attributed to the different tools used, which can lead to varied results. It is worth noting that our study relied on self-reported GPA, rather than obtaining the data directly from faculty administration.

In our study, students who reported extreme difficulty in achieving their work, managing household tasks, or interacting with others had significantly higher mean GAD scores. This finding is similar to previous studies that have reported a positive correlation between higher levels of anxiety severity and poorer quality of life,^{18,19} and lower health-related quality of life.⁴¹

Medical students with chronic diseases in our study exhibited higher mean GAD scores, consistent with earlier research that found significantly higher GAD scores among patients with conditions such as asthma,¹⁰ type 2 diabetes,⁹ and systemic lupus erythematosus (SLE).¹¹ These findings suggest that medical students with chronic diseases may be more susceptible to developing symptoms of GAD, which could have important implications for their mental health and overall well-being.

Surprisingly, our study demonstrates that medical students who engaged in regular exercise scored higher on the GAD-7 scale. This is contradictory to the existing literature, which suggests that physical activity plays an essential role in anxiety treatment⁴², particularly high-intensity exercise.^{43,44} However, this finding could be attributed to several factors. First, students with higher levels of anxiety might be more inclined to use regular exercise as a coping mechanism, leading to a correlation where those with higher anxiety levels are also those who exercise more. Second, medical students may engage in exercise to manage the high stress of their academic environment, which could contribute to higher anxiety levels despite regular physical activity. Additionally, it is noteworthy that the number of students who reported engaging in regular exercise was relatively small ($n=101$), which could influence the study's findings.

Strengths and Limitations

Our study provided significant insights into the mental health challenges faced by medical students in Sudan during a critical period. It was conducted at one of the largest and most prominent public universities in the country. However, the study

had certain limitations. While the GAD-7 tool was reliable, it's important to acknowledge that it functioned as a screening tool rather than a diagnostic instrument; a clinical assessment was necessary for a definitive diagnosis. The study was conducted at the Faculty of Medicine at Omdurman Islamic University. Although this is a large institution, this specific setting might have limited the ability to generalize our findings to medical students in other regions or countries. This suggested the need for multicenter studies. Furthermore, most of our study participants were female, which might have influenced the applicability of our findings across different genders. Moreover, the small sample size might have reduced the study's statistical power to detect significant effects.

Conclusion

This study revealed high GAD scores among Sudanese medical students, with significant associations found between GAD and female medical students as well as students with chronic diseases. GAD was found to negatively impact the ability of affected students to manage household tasks. The high prevalence of anxiety among medical students raises concerns, emphasizing the need for increased awareness, mainly focusing on recognizing and managing anxiety disorders. It is crucial to normalize discussions about mental health and provide easy access to counseling and psychotherapy services tailored to student needs. Additionally, fostering a culture of wellness within medical schools should be prioritized, with an emphasis on promoting healthy habits such as exercise, adequate sleep, and stress management. Furthermore, establishing peer support systems, including mentorship programs where senior students can guide and support juniors, can significantly contribute to a supportive environment. Additionally, training should be provided to faculty and administrative staff on how to identify and respond to students who experience anxiety, ensuring they can offer appropriate support or referrals. Future research should focus on evaluating the barriers to conducting and assessing the effectiveness of mental health services targeted at medical students and intervention programs aimed at reducing anxiety.

Summary – Accelerating Translation

Title: Understanding Anxiety among Medical Students in Sudan: A Study from Omdurman Islamic University

Main Problem: Anxiety, specifically Generalized Anxiety Disorder (GAD), was identified as a significant challenge globally, notably affecting the academic performance and overall quality of life of medical students. This problem was particularly pronounced in regions experiencing socio-political unrest, such as Sudan, where medical students were under increased stress due to the demanding nature of their studies compounded by external instability.

Aim: The study aimed to assess the prevalence of GAD among medical students at Omdurman Islamic University, identify the key factors that contributed to its development, and evaluate its impact on students' academic achievements and daily functioning. The goal was to provide insights that could aid in developing better support systems for medical

students' mental health, particularly in challenging environments like Sudan.

Methodology: This study was conducted among medical students at Omdurman Islamic University, a public university in Omdurman City, Sudan. The university was established in 1912 and has 22 faculties. In 2022, there were around 26,000 students. We used a method called systematic sampling to select participants for the study.

To collect data, we used an online questionnaire with two parts. The first part asked about things like age, gender, marital status, academic year, where the students lived, their grade point average, and medical history. The second part focused on generalized anxiety disorder (GAD) and used a questionnaire called the Generalized Anxiety Disorder Questionnaire (GAD-7) to measure its severity.

The study followed ethical guidelines and received approval from the Omdurman Islamic University Ethics Committee of the Faculty of Medicine. All participants gave their consent to participate in the study electronically.

Results: In this study, we looked at a total of 374 medical students who participated. It's interesting to note that most of them were female, making up 64.7% of the participants. Out of all the medical students in the study, 35 of them mentioned having a chronic disease. What's even more interesting is that more than one-third of these students, specifically 35.7%, reported having asthma.

Out of the 364 medical students who took part, 101 of them said they regularly exercise. It's worth mentioning that nearly half of these students, about 50%, mentioned doing cardio exercises on a regular basis. Additionally, around 40% of them said they regularly engage in walking exercises.

We found that about one-third of the medical students, specifically 33.7%, had high possibility of Generalized Anxiety Disorder (GAD). When we looked at the severity of anxiety, we found that 41.2% had mild anxiety, 21.4% had moderate anxiety, and 12.3% had severe anxiety.

To understand the differences in anxiety levels, we compared the average scores. It turned out that female medical students had significantly higher anxiety scores (average score of 7.80) compared to male medical students (average score of 6.06). However, we didn't find any significant differences in anxiety scores based on factors like age, city of residence, marital status, or living situation.

We found that fifth-year medical students are significantly more likely to experience anxiety than students in other academic years

Interestingly, medical students with chronic diseases had significantly higher anxiety scores (average score of 8.64) compared to those without chronic diseases (average score of 7.03). Moreover, we found a significant difference in anxiety scores between students who reported regular exercise (average score of 7.41) and those who didn't (average score of 6.29).

We also looked at the impact of anxiety on academic performance. However, we didn't find any significant association between anxiety and academic performance.

To understand how anxiety affects daily activities, we compared the average scores again. Students who reported extremely difficult daily activities had significantly higher anxiety scores (average score of 11.80), as did those who reported very difficult daily activities (average score of

11.95). On the other hand, students who found their daily activities not difficult (average score of 4.10) or somewhat difficult (average score of 6.86) had lower anxiety scores.

Conclusion: This study showed a high level of GAD among Sudanese medical students. Key insights include that high GAD is associated with female students and those suffering from chronic diseases and the negative effect of GAD on managing daily tasks. The concerning levels of anxiety among medical students underscore the urgent need for heightened awareness and better management strategies for anxiety

disorders. The study advocates for normalizing mental health discussions, ensuring accessible mental health services tailored to students, and promoting a wellness culture within medical educational institutions. Emphasizing healthy lifestyle choices, establishing supportive peer networks, and providing training for faculty and staff on recognizing and addressing student anxiety are pivotal steps. The conclusion calls for further research into the obstacles faced in implementing effective mental health services and interventions specifically designed to alleviate anxiety among medical students, aiming to enhance their overall well-being and academic success.

References

- World Health Organization. WHO highlights urgent need to transform mental health and mental health care. Available from: <https://www.who.int/news/item/17-06-2022-who-highlights-urgent-need-to-transform-mental-health-and-mental-health-care>. Last updated June 17, 2022; cited Feb 5, 2024.
- Centers for Disease Control and Prevention. Symptoms of Generalized Anxiety Disorder Among Adults: United States, 2019; Available from: <https://www.cdc.gov/nchs/products/index.htm>. Last updated Sep 2020; cited Feb 4, 2024
- Mishra AK, Varma AR. A Comprehensive Review of the Generalized Anxiety Disorder. *Cureus*. 2023;15(9):e46115.
- National Institute of Mental Health (NIMH). Generalized Anxiety Disorder. Available from: <https://www.nimh.nih.gov/health/statistics/generalized-anxiety-disorder>; cited Feb 4, 2024
- McLean CP, Asnaani A, Litz BT, Hofmann SG. Gender differences in anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness. *J Psychiatr Res*. 2011;45(8):1027–35.
- Bahrami F, Yousefi N. Females Are More Anxious Than Males: a Metacognitive Perspective. *Iran J Psychiatry Behav Sci*. 2011;5(2): 83–90.
- Vesga-López O, Schneier F, Wang S, Heimberg R, Liu SM, Hasin DS, et al. Gender differences in Generalized Anxiety Disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *J Clin Psychiatry*. 2008;69(10):1606–16.
- Gottschalk M, Domschke K. Genetics of generalized anxiety disorder and related traits. *Dialogues Clin Neurosci*. 2017;19(2):159–68.
- Huang CJ, Hsieh HM, Tu HP, Jiang HJ, Wang PW, Lin CH. Generalized anxiety disorder in type 2 diabetes mellitus: Prevalence and clinical characteristics. *Brazilian Journal of Psychiatry*. 2020;42(6):621–9.
- Lavoie KL, Boudreau M, Plourde A, Campbell TS, Bacon SL. Association between generalized anxiety disorder and asthma morbidity. *Psychosom Med*. 2011;73(6):504–13.
- Liao J, KJ, LF et al. A cross-sectional study on the Association of Anxiety and depression with the disease activity of systemic lupus erythematosus. *BMC Psychiatry*. 2022;22(1) 591.
- Quek TTC, Tam WWS, Tran BX, Zhang M, Zhang Z, Ho CSH, et al. The global prevalence of anxiety among medical students: A meta-analysis. *Int J Environ Res Public Health*. 2019;16(15):2735.
- Halperin SJ, Henderson MN, Prenner S, Grauer JN. Prevalence of Anxiety and Depression Among Medical Students During the Covid-19 Pandemic: A Cross-Sectional Study. *J Med Educ Curric Dev*. 2021;8:238212052199115.
- Wali RM, Bagabas TM, Hassanein AA, Saad Alameri M, Fahad K, Ouqla A. The Prevalence and Risk Factors of Depression and Anxiety Disorders among Medical Students in King Saud Bin Abdulaziz University for Health Sciences, Jeddah 2019. *Ann Med Health Sci Res*. 2021;11:1–10.
- Gabal HA, Wahdan MM, Gamal Eldin DA. Prevalence of anxiety, depression and stress among medical students, and associated factors. *Egypt J Occup Med*. 2022;46(1):55–74.
- Barbosa-Camacho FJ, Romero-Limón OM, Ibarrola-Peña JC, Almanza-Mena YL, Pintor-Belmontes KJ, Sánchez-López VA, et al. Depression, anxiety, and academic performance in COVID-19: a cross-sectional study. *BMC Psychiatry*. 2022 Dec;22(1)434.
- Jamil H, Alakkari M, Al-Mahini MS, Alsayid M, Al Jandali O. The Impact of Anxiety and Depression on Academic Performance: A Cross-Sectional Study among Medical Students in Syria. *Avicenna J Med*. 2022;12(03):111–9.
- Gan GG, Hue YL. Anxiety, depression and quality of life of medical students in Malaysia. *Med J Malaysia*. 2019;74(1):57–61.
- Wilmer MT, Anderson K, Reynolds M. Correlates of Quality of Life in Anxiety Disorders: Review of Recent Research. *Curr Psychiatry Rep*. 2021;23(11):77.
- Mohamed OGN, Mohamed EGN, Ahmed R, Aburas L, Ali M, Hamdan HZ. Depression, Anxiety, and Stress among Sudanese Medical Students during the COVID-19 Lockdown Period. *Open Access Maced J Med Sci*. 2022;10(B):1365–71.
- Mustafa SH, Mohammed EA, Makkawi ST, Mohammed YY. Mental Distress among Medical Students in Khartoum, Sudan 2022. *Open J Psychiatr*. 2022;12(04):345–58.
- Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006;166(10):1092–7.
- Löwe B, Decker O, Müller S, Brähler E, Schellberg D, Herzog W, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care*. 2008;46(3):266–74.
- Ibrahim D, Ahmed RM, Mohammad AZ, Ibrahim B, Mohammed T, Mohamed ME, et al. Prevalence and correlates of generalized anxiety disorder and perceived stress among Sudanese medical students. *BMC Psychiatry*. 2024;24(1):68.
- Alatawi A, Alghamdi A, Albalwi A, Altayar M, Jalal M, Frah EAM. Prevalence of Generalized Anxiety Disorder (GAD) Among Saudi Medical Students and Associated Risk Factors. *International Journal of Medical Research & Health Sciences*. 2020; 5(9):1–9
- Soltan MR, Soliman SS, Dawoud ME. A study of anxiety, depression and stress symptoms among Fayoum medical students during COVID-19 lockdown, Egypt. *Egypt J Neurol Psychiatr Neurosurg*. 2021;57(1):123.
- Fawzy M, Hamed SA. Prevalence of psychological stress, depression and anxiety among medical students in Egypt. *Psychiatry Res*. 2017; 255:186–94.
- Jafri SAM, Zaidi E, Aamir IS, et al. Stress Level Comparison of Medical and Nonmedical Students: A Cross Sectional Study done at Various Professional Colleges in Karachi, Pakistan. *Acta Psychopathologica*. 2017;03(02).
- Mohamed EY. Generalized anxiety disorder among saudi university medical students. *Academic Journal of Health Sciences* 2022;37(3):162–165.
- Çimen İD, Alvr TM, Coşkun B, Şükür NEÖ. Mental health of Turkish medical students during the COVID-19 pandemic. *Int J Soc Psychiatry*. 2022;68(6):1253–62.

31. Abdel Wahed WY, Hassan SK. Prevalence and associated factors of stress, anxiety and depression among medical Fayoum University students. *Alexandria Journal of Medicine*. 2017;53(1):77–84.
32. Seeman M V. Psychopathology in Women and Men: Focus on Female Hormones. *Am J Psychiatry*. 1997;154(12):1641–7.
33. Altemus M. Sex differences in depression and anxiety disorders: Potential biological determinants. *Horm Behav*. 2006;50(4):534–8.
34. Backović D V, Živojinović JI, Maksimović J, Maksimović M. Gender differences in academic stress and burnout among medical students in final years of education. *Psychiatr Danub*. 2012;24(2):175–81.
35. Villwock JA, Sobin LB, Koester LA, Harris TM. Impostor syndrome and burnout among American medical students: a pilot study. *Int J Med Educ*. 2016;7:364–9.
36. Rice J, Rosario-Williams B, Williams F, West-Livingston L, Savage D, Wilensky JA, et al. Impostor syndrome among minority medical students who are underrepresented in medicine. *J Natl Med Assoc*. 2023;115(2):191–8.
37. Niemi PM, Vainiomäki PT. Medical students' distress - Quality, continuity and gender differences during a six-year medical programme. *Med Teach*. 2006;28(2):136–41.
38. Dyrbye LN, Thomas MR, Shanafelt TD. Medical student distress: Causes, consequences, and proposed solutions. *Mayo Clin Proc*. 2005;80(12):1613–22.
39. von der Embse N, Jester D, Roy D, Post J. Test anxiety effects, predictors, and correlates: A 30-year meta-analytic review. *J Affect Disord*. 2018;227:483–493.
40. Teh CK, Ngo CW, Zulkifli RA binti, Vellasamy R, Suresh K. Depression, Anxiety and Stress among Undergraduate Students: A Cross Sectional Study. *Open J Epidemiol*. 2015;05(04):260–8.
41. Comer JS, Blanco C, Hasin DS, Liu SM, Grant BF, Turner JB, et al. Health-related quality of life across the anxiety disorders: Results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Journal of Clinical Psychiatry*. 2011;72(1):43–50.
42. Mc Dowell CP, Carlin A, Capranica L, Dillon C, Harrington JM, Lakerveld J, et al. Associations of self-reported physical activity and anxiety symptoms and status among 7,874 Irish adults across harmonised datasets: A DEDIPAC-study. *BMC Public Health*. 2020 20;20(1).
43. Aylett E, Small N, Bower P. Exercise in the treatment of clinical anxiety in general practice - A systematic review and meta-analysis. *BMC Health Serv Res*. 2018 ;18(1):559.
44. Herring MP, Monroe DC, Gordon BR, Hallgren M, Campbell MJ. Acute Exercise Effects among Young Adults with Analogue Generalized Anxiety Disorder. *Med Sci Sports Exerc*. 2019;51(5):962–9.

Acknowledgments

We would like to express our appreciation to the invited reviewers for their valuable contribution, which enhanced the quality of the work. We also extend our gratitude to Wafa, Muna, and Maab for their tireless work as data collectors during the data gathering phase.

Conflict of Interest Statement & Funding

The Authors have no funding, financial relationships or conflicts of interest to disclose.

Author Contributions

K.O.M, E.A.Z, W.A.A, and A.M.M. Methodology: K.O.M. Formal Analysis: K.O.M. Writing-Original Draft: K.O.M, A.A.A, E.A.Z, S.M.S, W.A.A, and A.M.M. Writing-Review & Editing: K.O.M, A.A.A, E.A.Z, and S.M.S. Visualization: A.A.A. Project Administration: K.O.M.

Cite as

Mohamed KO, Ahmed AA, Zaki EA, Soumit SM, Allam WA, Mohamed AM. Prevalence of Generalized Anxiety Disorder and Associated Risk Factors Among Medical Students in Sudan: A Cross-Sectional Study at Omdurman Islamic University. *Int J Med Stud*. 2024 Jan-Mar;12(1):14–21.

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

ISSN 2076-6327

This journal is published by [Pitt Open Library Publishing](https://open.library.pitt.edu/)

