

1 **Title: A Cross Sectional Study on Adherence to Medication among Patients with Hypertension and/or**
2 **Diabetes Attending One of the Tertiary Care Institutes of Ahmedabad City, Gujarat, India**

3
4 **Article type:** Original Article

5
6 **Author names:**

- 7 1. Janak Patel
- 8 2. Kanal Shah
- 9 3. Jayveer Jain
- 10 4. Venu Shah

11
12 **Degrees and Affiliations:**

- 13 1. Third year Medical Student. GCS Medical College, Hospital & Research Center, Ahmedabad, India.
- 14 2. Third year Medical Student. GCS Medical College, Hospital & Research Center, Ahmedabad, India.
- 15 3. Third year Medical Student. GCS Medical College, Hospital & Research Center, Ahmedabad, India.
- 16 4. MD Community Medicine. Professor, Community Medicine Department, GCS Medical College, Hospital
17 & Research Center, Ahmedabad, India.

18
19 **ORCID (Open Researcher and Contributor Identifier):**

20 <https://orcid.org/0009-0006-0315-8585>

21 <https://orcid.org/0009-0005-2330-9769>

22 <https://orcid.org/0009-0005-8822-3034>

23 <https://orcid.org/0000-0003-4133-582X>

24
25 **Corresponding author email:** drvenushah@gmail.com

26 **Financing:** Nil

27 **Conflict of interest statement by authors:** Nil

28
29 **Authors Contribution Statement:** Conceptualization: JP, KS, JJ, VS. Data Curation: JP, KS, JJ. Formal
30 Analysis: JP, KS, JJ, VS. Investigation: KS, JJ. Methodology: JP, KS, JJ, VS. Supervision: VS. Validation:
31 VS. Writing - Original Draft: JP, KS, JJ, VS. Writing - Review Editing: VS.

32
33 **Manuscript word count:** 1690

34 **Abstract word count:** 242

35 **Number of Figures and Tables:** 4 (Tables)

1 **ABSTRACT:**

2 **Background:** The lifelong management of chronic diseases such as Hypertension and Diabetes Mellitus
3 necessitates a comprehensive approach, including lifestyle modifications and consistent adherence to
4 medication. Present study aimed to evaluate treatment adherence among patients with diabetes and /or
5 hypertension attending a tertiary care institute in Ahmedabad city, India.

6 **Methods:** A cross-sectional study was conducted among 200 patients aged over 18 years, diagnosed with
7 diabetes and/or hypertension. The sample population was selected from a tertiary care institute at Ahmedabad
8 city using a consecutive sampling method. Data collection utilized a pretested and predesigned proforma along
9 with the Morisky Medication Adherence Scale (MMAS-8).

10 **Results:** Hypertension was present in 43.0% of the participants, while 20.0% reported diabetes mellitus.
11 Additionally, 37.0% had both conditions. Out of total 200 patients 53.5% exhibited low adherence to treatment.
12 Lower adherence to treatment was found to be more among Diabetics (67.5%) as compared to hypertensive
13 patients (51.2%). Among participants having comorbidities 48.6% had low adherence to treatment. The
14 association between disease type and adherence levels was not significant (p-value = 0.305). Not having any
15 symptoms and forgetfulness were common reasons for low adherence among 41.1% and 24.2% of participants,
16 respectively.

17 **Conclusion:** This study highlights the need for targeted interventions aimed at enhancing medication
18 adherence among individuals managing with chronic illnesses like hypertension and diabetes. The findings
19 provide valuable insights for healthcare professionals and policymakers to develop effective strategies for
20 improving treatment adherence and subsequently mitigating the progression of these chronic conditions.

21
22 **Key Words:** Morisky Medication Adherence, Hypertension, Diabetes

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

1 Dates

2 Submission: 03/09/2024

3 Revisions: 07/18/2024, 09/16/2024, 09/26/2024

4 Responses: 07/24/2024, 09/23/2024, 10/04/2024

5 Acceptance: 10/25/2024

6 Publication: 11/18/2024

7

8 Editors

9 Associate Editor/Editor: Francisco J. Bonilla-Escobar

10 Student Editors: Prajjwol Luitel, Siddhant Govekar & Mirna Mustafa

11 Copyeditor: Emmanuel Phiri

12 Proofreader:

13 Layout Editor:

14

15 **Publisher's Disclosure:** *This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our readers and authors we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.*

20

21

22

23 INTRODUCTION:

24 Non-communicable diseases are on the brink of a surge in developing countries. India, already
25 acknowledged as the "Diabetic capital of the world," is now progressing toward earning the title of the
26 "Hypertension capital of the world" as well.¹ Lifelong treatment is mandatory in both the diseases to avert any
27 complications. Adherence to antihypertensive as well as anti-diabetic treatment has a key role in effective
28 management of the respective disease conditions. The World Health Organization (WHO) defined adherence
29 as the extent to which a patient follows medical instructions.² In India, studies have reported rates of non-
30 adherence to medication among the hypertensive and diabetic patients to be between 25 % and 85%.³⁻⁶
31 Factors affecting adherence to medication among hypertensive and diabetic patients were identified in
32 previous studies which included type of drug regimen, duration of treatment as well as psychosocial factors.⁷

33 Despite the growing burden of diabetes and hypertension in India, there is a scarcity of
34 comprehensive studies addressing adherence to treatment in the current study setting. Recognizing this
35 research gap, the present study intended to assess medication adherence among patients with diabetes and
36 hypertension attending one of the tertiary care hospitals in Ahmedabad city. It was also envisioned to find out
37 the reasons behind low adherence to the treatment pertaining to diabetes and/or hypertension. This
38 investigation is crucial for informing healthcare practices, enhancing treatment strategies, and ultimately
39 improving the overall health outcomes of individuals grappling with these chronic conditions.

40

41 METHODS:

1 Cross-sectional study was conducted among patients attending Out patients Department of tertiary care
2 teaching institute of Ahmedabad city of India. Study participants were patients already diagnosed with
3 Hypertension and/or Diabetes taking treatment for more than 6 months.

4 Inclusion criteria: Any case of hypertension and/or diabetes more than 18 years of age diagnosed with
5 diabetes and/or hypertension for more than 6 months ready to give consent for the study.

6 Exclusion criteria: Patients with any psychiatric illness not able to respond to questionnaire and Diabetic
7 patients taking insulin injections were excluded from the study.

8 Sample size was calculated using the formula $4pq / l^2$, where p is the adherence to treatment among
9 diabetic patients as per previous study i.e.83.6% ⁸absolute precision of 5.5%, power of 80%, at 95% confidence
10 and 10 % of non response rate, the sample size calculated to 192, which was rounded up to 200.Using
11 consecutive sampling, patients attending medical OPD of tertiary care during study period were interviewed till
12 the sample size of 200 was achieved.

13 Personal interview of the study participants was conducted using pre-tested and pre-designed
14 proforma. Main domains of the questionnaire include: Socio demographic details, Medical history related to
15 diabetes and / or hypertension, nature of treatment, personal history etc. Socioeconomic status was assessed
16 using modified Prasad scale.⁹Adherence to treatment was determined by Morisky medication adherence scale
17 MMAS-8.¹⁰ In this scale, the first seven questions had binary response categories (yes/no) while the eighth item
18 had a five-point Likert response. The scores of the MMAS-8 range from 0 to 8. A score below 6 indicates low
19 adherence, a score between 6 < 8 medium adherence and a score of 8 high adherence. Previous study¹¹
20 suggested that MMAS-8 had better psychometric properties, with sensitivity and specificity of 93% and 53%
21 respectively and Cronbach's alpha value of 0.83.

22 Data was entered in Micro Soft Excel and analysed using Jamovi Software. The categorical variables
23 expressed in frequency and percentages and the association between the two groups was calculated using
24 Chi-square test. The quantitative variables were expressed as mean (standard deviation). A p-value of less than
25 0.05 was considered as statistically significant.

26

1 **RESULTS:**

2 The majority of the study participants falls within the age group of 46-60 years, comprising 46.5% of the study
3 population. There was a near-equal distribution of gender, with females at 48.0% and males at 52.0%. The
4 most common education level among study group was secondary (32.5%), followed by graduate (20.5%). A
5 significant portion of the population was unemployed (56.0%) and belonged to joint families (56.0%).
6 Maximum numbers of participants were from upper socio-economic class constituting 47.0% of the sample.
7 (Table 1)

8 Among total participants, Hypertension was the most prevalent condition affecting 43.0% of the
9 sample where as Diabetes mellitus was reported by 20.0% of individuals. Total 37.0% of the respondents
10 were having with both hypertension and diabetes mellitus. A significant portion of study participants (52.0%)
11 were diagnosed with above condition for more than 5 years. Approximately 19.5% each fall into the categories
12 of 1-3 years, and 3-5 years since diagnosis. The majorities (72.0%) were taking less than 3 pills per day and
13 only 5.5% took more than 5 pills daily. Out of pocket expenditure pertaining to medication expenses was
14 assessed. It was found that 29.0% participants were spending between 200-400 INR on medicines while
15 27.0% had expense of more than 600 INR. Majority (83.5%) participants were adhered to regular follow-up
16 visits as well as lifestyle modification after diagnosis of disease. (Table 2)

17 Overall around 107 (53.5%) participants were having low adherence to treatment. For diabetes
18 mellitus, a higher percentage of individuals exhibited low adherence (67.5%), with none reported high
19 adherence. In the case of hypertension, low adherence was seen in 51.2% cases, while 3.5% had high
20 adherence to treatment. Individuals with both diabetes mellitus and hypertension had a balanced distribution
21 (48.6% each) between low and medium adherence, with a smaller percentage showing high adherence
22 (2.7%). The Chi-square test was applied to evaluate the association between disease type and adherence
23 levels. However, it was not statistically significant (p -value = 0.305). (Table 3)

24 On assessment of possible reasons behind low adherence, it was found that a significant portion of
25 individuals (41.1%) were not adhered to their treatment plan because they perceived no signs or symptoms,
26 potentially underestimating the importance of continued medication. Forgetfulness was also a common reason
27 for low adherence among 24.2% participants. (Table 4)

28
29
30
31

DISCUSSION:

Current study evaluated the adherence to treatment among patients with Diabetes, Hypertension as well as patients having both of these comorbidities. Total 200 patients were interviewed using pretested proforma including morisky medication adherence scale. The study population predominantly comprised individuals aged 46-60 years (46.5%), with a balanced gender distribution (48.0% females, 52.0% males). Secondary education was the most common (32.5%), followed by graduate (20.5%), and majority belonged to joint families (56.0%). Hypertension affected 43.0%, diabetes mellitus 20.0%, and 37.0% had both conditions. A significant portion (52.0%) had been diagnosed for over 5 years. Most participants (72.0%) consumed less than 3 pills daily, with 29.0% spending 200-400 INR on medicines. The majority (83.5%) adhered to follow-up visits and lifestyle modifications.

Patients diagnosed with Diabetes Mellitus, 67.5% demonstrated low adherence to treatment, while among those diagnosed with Hypertension, 51.2% exhibited low adherence. Those having both the illnesses, 48.6% had low adherence to treatment. In the study conducted by Balasubramanian et al among hypertensive patients of rural Kerala, 41.3% had high adherence, while the remaining 58.7% had either medium or low adherence.¹² Treatment adherence for hypertension was 82% in study conducted by Rao et al.¹³ One of the meta-analysis conducted by Pokharel et al reported that half of the hypertensive population of Nepal were non-adherent to their anti-hypertensive medications.¹⁴

In current study, 67.5% of diabetic patients reported low adherence to treatment. Similar finding were observed in the study conducted by Sahoo et al with 65.8% of diabetic patients attending tertiary care hospital of Bhubaneswar, India with poor adherence to treatment.¹⁵ Contrary to these findings Rao et al reported in their study that 83.6% diabetic patients were compliant to treatment.¹³ In current study hypertensive patients had better adherence to treatment as compared to Diabetic patients, whereas Sweileh et al noted that diabetic patients had better overall rate of compliance than hypertensive patients in their study.¹⁶ Difference in adherence rates observed in various studies could be due to variation in the methods utilized for assessment of adherence. Although adherence levels varied for diabetes mellitus and hypertension, the association between disease type and adherence was not statistically significant in the current study (p-value = 0.305). It's important to note that while these statistics provide insights into the relationship between disease type and adherence, a non-significant result does not imply the absence of a relationship as it may be influenced by the sample size.

Current study evaluated the reasons behind low adherence among the study participants. Lack of symptoms perception (41.1%) and forgetfulness (24.2%) were the commonest reasons mentioned by the study cohort. Similar finding was present in the study conducted by Pokharel et al who mentioned forgetfulness, carelessness and cost of medications as the primary reasons for non-adherence to treatment.¹⁴ Rao et al identified various factors contributing to low treatment adherence, with the most prevalent reasons including the cost of medications and the asymptomatic nature of the disease.¹³ Lack of symptoms in the disease can impede the patient's motivation for treatment since there are no tangible benefits experienced by the patient.

The current study concluded that overall treatment adherence was low among the study population affecting with diabetes and/or hypertension. The study highlights the need for targeted interventions to address low adherence, considering factors such as perception and forgetfulness. This research contributes to the existing literature on adherence to medication in chronic diseases, offering a basis for future interventions and public health initiatives.

SUMMARY - ACCELERATING TRANSLATION:

1 A Cross Sectional Study was conducted on Adherence to Medication among Patients with Hypertension
2 and/or Diabetes Attending One of the Tertiary Care Institutes of Ahmedabad City of India. Total 200 patients
3 were interviewed in the study to identify level of adherence to treatment among them. Pretested and
4 predesigned proforma including Morisky medication Adherence scale was used for data collection. It was
5 observed that patients having Diabetes, Hypertension as well as patients with both of these conditions were
6 having poor adherence to treatment with percentage of low adherence 67.5, 51.2 and 48.6%, respectively. On
7 evaluating the reasons for low adherence it was found that patients were ignoring the disease condition due to
8 absence of any symptoms. Forgetfulness in taking medications was also mentioned by one fourth of patients.
9 The study highlights the need for targeted intervention towards increasing adherence to medication among
10 patients with chronic illnesses like hypertension and diabetes.

11 REFERENCES:

- 12 1. Joshi SR, Parikh RM. India – Diabetes capital of the world: Now heading towards hypertension. J Assoc
13 Physicians India. 2007;55:323–4.
- 14 2. World Health Organization (WHO). What is adherence. WHO.2003. pages 17–18. Available from:
15 https://www.who.int/chp/knowledge/publications/adherence_Section1.pdf. Cited on May 29 2023
- 16 3. Bhandari S, Sarma PS, Thankappan KR. Adherence to antihypertensive treatment and its determinants
17 among urban slum dwellers in Kolkata, India. Asia Pac J Public Health.
- 18 4. Venkatachalam J, Abrahm SB, Singh Z, Stalin P, Sathya GR. Determinants of Patient's Adherence to
19 Hypertension Medications in a Rural Population of Kancheepuram District in Tamil Nadu, South India.
20 Indian J Community Med. 2015 Jan-Mar;40(1):33-7.
- 21 5. Mishra R, Sharma SK, Verma R, Kangra P, Dahiya P, Kumari P, Sahu P, Bhakar P, Kumawat R, Kaur
22 R, Kaur R, Kant R. Medication adherence and quality of life among type-2 diabetes mellitus patients in
23 India. World J Diabetes. 2021 Oct 15;12(10):1740-1749.
- 24 6. Venkatesan M, Dongre AR, Ganapathy K. A Community-Based Study on Diabetes Medication
25 Nonadherence and its Risk Factors in Rural Tamil Nadu. Indian J Community Med. 2018 Apr-
26 Jun;43(2):72-76.
- 27 7. Lavakumar S, Jesurun RS. A study on the level of drug compliance among the outpatients who are on
28 a long-term drug therapy in a tertiary care teaching hospital at Kancheepuram district in Tamil Nadu.
29 Asian J Pharm Clin Res 2017;10:174-6.
- 30 8. Rao CR, Kamath VG, Shetty A, Kamath A. Treatment Compliance among Patients with Hypertension
31 and Type 2 Diabetes Mellitus in a Coastal Population of Southern India. Int J Prev Med. 2014
32 Aug;5(8):992-8.
- 33 9. Pentapati SSK, Debnath DJ. Updated BG Prasad's classification for the year 2022. J Family Med Prim
34 Care. 2023 Jan;12(1):189-190.
- 35 10. Morisky DE, Ang A, Krousel-Wood M, Ward HJ. Predictive validity of a medication adherence measure
36 in an outpatient setting. J Clin Hypertens (Greenwich). 2008 May;10(5):348-54.
- 37 11. Surekha A, Fathima FN, Agrawal T, Misquith D. Psychometric properties of Morisky Medication
38 Adherence Scale (MMAS) in known diabetic and hypertensive patients in a rural population of Kolar
39 District, Karnataka. Indian J Public Heal Res Dev. 2016;7:250–256.
- 40 12. Balasubramanian A, Nair SS, Rakesh PS, Leelamoni K. Adherence to treatment among hypertensives
41 of rural Kerala, India. J Family Med Prim Care. 2018 Jan-Feb;7(1):64-69.

- 1 13. Rao CR, Kamath VG, Shetty A, Kamath A. Treatment Compliance among Patients with Hypertension
2 and Type 2 Diabetes Mellitus in a Coastal Population of Southern India. Int J Prev Med. 2014
3 Aug;5(8):992-8.
- 4 14. Pokharel P, Jha SK, Adhikari A, Katwal S, Ghimire S, Shrestha AB, Poudel N. Non-adherence to anti-
5 hypertensive medications in a low-resource country Nepal: a systematic review and meta-analysis. Ann
6 Med Surg (Lond). 2023 Jul 19;85(9):4520-4530.
- 7 15. Sahoo J, Mohanty S, Kundu A, Epari V. Medication Adherence Among Patients of Type II Diabetes
8 Mellitus and Its Associated Risk Factors: A Cross-Sectional Study in a Tertiary Care Hospital of Eastern
9 India. Cureus. 2022 Dec 29;14(12):e33074.
- 10 16. Sweileh WM, Aker O, Hamooz S. Rate of compliance among patients with diabetes mellitus and
11 hypertension. An-Najah Univ J Research-A-(Natural Sciences) 2005;19:1–12.

Accepted, in-press

37 **Table 1: Demographic profile of study participants (n=200)**

Variables	n	%
Age Group (Years)		
18 - 30	4	2.0

31 - 45	27	13.5
46 - 60	93	46.5
61 - 75	65	32.5
More than 75	11	5.5
Gender	5	
Female	96	48.0
Male	104	52.0
Education		
Primary	31	15.5
Secondary	65	32.5
Higher secondary	36	18.0
Graduate	41	20.5
Post graduate	27	13.5
Occupation		
Employed	88	44.0
Unemployed	112	56.0
Type of family		
Joint	112	56.0
Nuclear	81	40.5
Three generations	7	3.5
Socio Economic Class		
Upper	94	47.0
Upper Middle	57	28.5
Middle	24	12.0
Lower Middle	13	6.5
Lower	12	6.0

7
8
9
10
11
12
13
14
15
16
17
18

Table 2: Profile related to disease and its management among study group (n=200)

Variables	n	%
Type of disease		

Hypertension	86	43.0
Diabetes mellitus	40	20.0
Both	74	37.0
Duration since diagnosis of disease		
Less than 1 year	18	9.0
1-3 year	39	19.5
3-5 year	39	19.5
More than 5 year	104	52.0
Number of pills per day		
Less than 3	144	72.0
3 to 5	45	22.5
More than 5	11	5.5
Expense incurred for purchase of medicine (INR)		
<200	44	22.0
200-400	58	29.0
400-600	44	22.0
>600	54	27.0
Regular Follow up visit		
Yes	167	83.5
No	33	16.5
Life style modification after diagnosis of disease		
Yes	167	83.5
No	33	16.5
Total	200	100

1
2
3
4
5
6
7
8
9
10
11
12
13

Table 3: Adherence to treatment among patients (n=200)

Disease	Adherence to Treatment			Chi square (p value)
	Low	Medium	High	

Diabetes mellitus (n=40)	27 (67.5 %)	13 (32.5 %)	0 (0.0 %)	4.83 (0.305)
Hypertension (n=86)	44(51.2 %)	39 (45.3 %)	3 (3.5 %)	
Both (n=74)	36 (48.6%)	36 (48.6 %)	2(2.7 %)	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

Accepted, in-press

Table 4: Reasons mentioned by participant for low adherence to treatment (n=107)

Reasons for low adherence*	n	%
----------------------------	---	---

No signs and symptoms	44	41.1
Forgetfulness	26	24.2
Non availability of medicine	22	20.5
High cost of medicines	19	17.7
Side effects	12	11.2

1 *Multiple responses

Accepted, in-press